arXiv:1206.5685v4 [hep-ph] 25 Mar 2013

Radiative transitions among the vector and scalar heavy quarkonium states with covariant light-front quark model

Zhi-Gang Wang [1]
Department of Physics, North China Electric Power University, Baoding 071003, P. R. China

Abstract

In this article, we study the radiative transitions among the vector and scalar heavy quarkonium states with the covariant light-front quark model. In calculations, we observe that the radiative decay widths are sensitive to the constituent quark masses and the shape parameters of the wave-functions, and reproduce the experimental data with suitable parameters.

PACS numbers: 14.40.Pq; 13.40.Hq
Key Words: Covariant light-front quark model, Radiative decays

1 Introduction

Recently, the BESIII collaboration observed the first evidence for direct two-photon transition $\psi' \to J/\psi\gamma\gamma$ with the branching fraction $(3.3 \pm 0.6^{+0.8}_{-1.1}) \times 10^{-4}$ in a sample of 106 million ψ' decays collected by the BESIII detector [1]. The branching fractions of the double E_1 transitions $\psi' \to (J/\psi\gamma)_{\chi_{cj}}\gamma$ through the intermediate states χ_{cj} $(j = 0, 1, 2)$ are also reported [1], while previous experimental data indicates that the double radiative decays $\psi' \to J/\psi\gamma\gamma$ take place through the decay cascades $\psi' \to \chi_{cj}\gamma \to J/\psi\gamma\gamma$ with tiny non-resonance contributions [2, 3]. In Ref.[4], He et al study the discrete contributions to decays $\psi' \to J/\psi\gamma\gamma$ due to the E_1 transitions using the heavy quarkonium effective Lagrangian [5]. No theoretical work on the non-resonance's contributions exists. On the other hand, we expect that there are non-resonance's contributions to the doubly radiative decays among the bottomnium states $\Upsilon(nS)$, the doubly radiative transition $\Upsilon'' \to \Upsilon'\gamma\gamma$ has been observed [6].

The radiative transitions among the heavy quarkonium states are usually calculated model-dependently by the nonrelativistic potential quark models with considerable relativistic corrections [7], or calculated model-independently by the lattice QCD [8] and effective field theory [9], one can consult the recent review [10] for more references. In general, we expect to study both the resonance's and non-resonance's contributions in the doubly radiative transitions with the covariant light-front quark model, where the wave-functions are expressed in terms of the internal variables of the quark and gluon, and maintain Lorentz covariance. In Ref.[11], Jaus introduces the covariant light-front quark model and takes into account the zero-mode contributions systematically to preserve covariance and remove dependence of physical quantities on the light-front direction. The covariant light-front quark model has been successfully applied to calculate the S-wave and P-wave meson's decay constants and form-factors [12, 13, 14]. In Refs.[15, 16], the authors study the M_1 transitions $\Upsilon(nS) \to \eta_b(n'S)\,\gamma$ in the covariant light-front quark model, and observe that the M_1 transitions are sensitive to the heavy quark masses and shape parameters of the wave-functions, the existing experimental data cannot be reproduced consistently with suitable parameters [15]. It is interesting to study whether or not the covariant light-front quark model can be successfully applied to calculate the E_1 transitions among the heavy quarkonium states.

In this article, we intend to study the radiative transitions (the E_1 transitions) among the vector and scalar heavy quarkonium states using the covariant light-front quark model as the first step, because they can be taken as the sub-processes of the doubly radiative decays, and the calculations are relatively simple. Experimentally, the widths of the radiative decays $\psi' \to \chi_{c0}\,\gamma$, $\chi_{c0} \to J/\psi\,\gamma$, $\Upsilon' \to \chi_{b0}\,\gamma$, $\Upsilon'' \to \chi_{b0}\,\gamma$, $\Upsilon'' \to \chi'_{b0}\,\gamma$ have been precisely measured [6]. A large amount of bottomonium states will be produced at the Large Hadron Collider, and the radiative transitions will be studied experimentally. We can study the radiative transitions by measuring

---

[1] E-mail:wangzgyiti@yahoo.com.cn.



Figure 1: The Feynman diagrams contribute to the form-factors. The photon is emitted from the quark (antiquark) line in the diagram A (B).

the energy spectrum of the photons or reconstructing the final quarkonium states, although the soft photons are difficult to identify. The four-photon decay cascades $\Upsilon(3S) \to \gamma\chi_{bj}(2P) \to \gamma\gamma\Upsilon(1D) \to \gamma\gamma\gamma\chi_{bj}(1P) \to \gamma\gamma\gamma\gamma\Upsilon(1S) \to \gamma\gamma\gamma\gamma\ell^+\ell^-$ ($j = 0, 1, 2$) have been observed by the CLEO collaboration [17], where the softest photons have the energy less than $90\,\mathrm{MeV}$.

The article is arranged as follows: we calculate the radiative transitions among the vector and scalar heavy quarkonium states with the covariant light-front quark model in Sect.2; in Sect.3, we present the numerical results and discussions; and Sect.4 is reserved for our conclusions.

2 Radiative decays with covariant light-front quark model

The radiative transitions among the heavy quarkonium states can be described by the following electromagnetic lagrangian $\mathcal{L}$,

$$\mathcal{L} = -ee_b\bar{b}\gamma_\mu bA^\mu - ee_c\bar{c}\gamma_\mu cA^\mu \, , \tag{1}$$

where the A_μ is the electromagnetic field, the e is the electromagnetic coupling constant, $e_b = -\frac{1}{3}$, and $e_c = \frac{2}{3}$. The transition amplitudes $\xi^\mu\mathcal{T}_\mu^{V\to S\gamma}$ and $\xi^\mu\mathcal{A}_\mu^{S\to V\gamma}$ can be decomposed as

$$\begin{aligned}\mathcal{T}_\mu^{V\to S\gamma} &= f_{VS\gamma}\left(\epsilon_\mu q\cdot P' - P'_\mu\epsilon\cdot q\right) \, , \\ \mathcal{A}_\mu^{S\to V\gamma} &= g_{SV\gamma}\left(\epsilon^*_\mu q\cdot P - P_\mu\epsilon^*\cdot q\right) \, , \end{aligned} \tag{2}$$

according to Lorentz covariance, where the V and S denote the vector and scalar mesons respectively, the P, P', q are the momenta of the initial mesons, final mesons and photons respectively, the ξ_μ and ϵ_μ are the polarization vectors of the photons and vector mesons respectively, the $f_{VS\gamma}$ and $g_{SV\gamma}$ are the electromagnetic form-factors (or the coupling constants) at $q^2 = 0$. The amplitudes $\mathcal{T}_\mu^{V\to S\gamma}$ and $\mathcal{A}_\mu^{S\to V\gamma}$ satisfy conservation of electromagnetic currents, we can replace the photon's polarization vector ξ_μ with its momentum q_μ in the transition amplitudes $\xi^\mu\mathcal{T}_\mu^{V\to S\gamma}$ and $\xi^\mu\mathcal{A}_\mu^{S\to V\gamma}$ to obtain $q^\mu\mathcal{T}_\mu^{V\to S\gamma} = q^\mu\mathcal{A}_\mu^{S\to V\gamma} = 0$.

From the lagrangian $\mathcal{L}$, we can draw the Feynman diagrams (see Fig.1) and write down the transition amplitudes,

$$\mathcal{T}_\mu^{V\to S\gamma} = -iee_QN_c\int\frac{d^4p_1}{(2\pi)^4}\left[\frac{H_VH_S}{N_1N_2N'_1}s^A_{\mu\alpha} + \frac{H_VH_S}{N_1N_2N'_2}s^B_{\mu\alpha}\right]\epsilon^\alpha_V \, , \tag{3}$$

$$\mathcal{A}_\mu^{S\to V\gamma} = -iee_QN_c\int\frac{d^4p_1}{(2\pi)^4}\left[\frac{H_SH_V}{N_1N_2N'_1}\widetilde{s}^A_{\mu\alpha} + \frac{H_SH_V}{N_1N_2N'_2}\widetilde{s}^B_{\mu\alpha}\right]\epsilon^{*\alpha}_V \, , \tag{4}$$

where $N_i = p_i^2 - m_Q^2$, $N_i' = p_i'^2 - m_Q^2$, and

$$\begin{aligned}
s^A_{\mu\alpha} &= \mathrm{Tr}\left\{(\not{p}_1' + m_Q)\gamma_\mu(\not{p}_1 + m_Q)\left[\gamma_\alpha - \frac{(p_1-p_2)_\alpha}{W_V}\right](-\not{p}_2 + m_Q)\right\}\,, \\
s^B_{\mu\alpha} &= \mathrm{Tr}\left\{(\not{p}_1 + m_Q)\left[\gamma_\alpha - \frac{(p_1-p_2)_\alpha}{W_V}\right](-\not{p}_2 + m_Q)\gamma_\mu(-\not{p}_2' + m_Q)\right\}\,, \qquad (5)\\
\widetilde{s}^A_{\mu\alpha} &= \mathrm{Tr}\left\{(\not{p}_1' + m_Q)\gamma_\mu(\not{p}_1 + m_Q)(-\not{p}_2 + m_Q)\left[\gamma_\alpha - \frac{(p_1'-p_2)_\alpha}{W_V}\right]\right\}\,, \\
\widetilde{s}^B_{\mu\alpha} &= \mathrm{Tr}\left\{(\not{p}_1 + m_Q)(-\not{p}_2 + m_Q)\gamma_\mu(-\not{p}_2' + m_Q)\left[\gamma_\alpha - \frac{(p_1-p_2')_\alpha}{W_V}\right]\right\}\,, \qquad (6)
\end{aligned}$$

the H_V and H_S are the light-front wave-functions of the vector and scalar mesons respectively, the A and B denote the diagrams in which the photon emitted from the quark and antiquark lines, respectively (see Fig.1), and the W_V is a parameter. In writing the transition amplitudes $\mathcal{T}_\mu^{V\to S\gamma}$ and $\mathcal{A}_\mu^{S\to V\gamma}$, we have used the following definitions of the quark-meson-antiquark vertexes $i\Gamma_M$ [12],

$$\begin{aligned}
i\Gamma_M &= -iH_S && \text{for scalar meson}\,, \\
&= iH_V\left[\gamma_\mu - \frac{(p_1-p_2)_\mu}{W_V}\right] && \text{for vector meson}\,, \qquad (7)
\end{aligned}$$

where the p_1 and p_2 are momenta of the quark and antiquark, respectively.

We assume that the covariant light-front wave-functions H_V and H_S are analytic in the upper (lower) p_1^- complex plane, close the integral contour in the upper (lower) p_1^- complex plane for the diagram A (B), which corresponds to set the antiquark (quark) on the mass-shell. The one-shell restrictions are implemented by the following replacements,

$$\begin{aligned}
p_2 &\to \hat{p}_2^2 = m_Q^2\,,\ \ p_1 \to \hat{p}_1 = P - \hat{p}_2\,, \\
N_1 &\to \hat{N}_1 = \hat{p}_1^2 - m_Q^2\,,\ \ N_1' \to \hat{N}_1' = \hat{p}_1'^2 - m_Q^2\,, \\
H_V &\to \hat{H}_V = h_V(x_2,p_\perp)\,,\ \ H_S \to \hat{H}_S = h_S(x_2,p_\perp)\,,\ \ W_V \to \hat{W}_V = w_V\,, \\
\int \frac{d^4p_1}{N_1N_2N_1'} &\to -i\pi\int\frac{dx_2d^2p_\perp}{x_2\hat{N}_1\hat{N}_1'}\,, \qquad (8)
\end{aligned}$$

for the Feynman diagram A and

$$\begin{aligned}
p_1 &\to \hat{p}_1^2 = m_Q^2\,,\ \ p_2 \to \hat{p}_2 = P - \hat{p}_1\,, \\
N_2 &\to \hat{N}_2 = \hat{p}_2^2 - m_Q^2\,,\ \ N_2' \to \hat{N}_2' = \hat{p}_2'^2 - m_Q^2\,, \\
H_V &\to \hat{H}_V = h_V(x_2,p_\perp)\,,\ \ H_S \to \hat{H}_S = h_S(x_2,p_\perp)\,,\ \ W_V \to \hat{W}_V = w_V\,, \\
\int \frac{d^4p_1}{N_1N_2N_2'} &\to -i\pi\int\frac{dx_2d^2p_\perp}{x_1\hat{N}_2\hat{N}_2'}\,, \qquad (9)
\end{aligned}$$

for the Feynman diagram B, where we have used the light-front decomposition of the momenta $P = (P^+, P^-, 0_\perp)$, $P^\pm = P^0 \pm P^3$, $p_{1,2}^+ = x_{1,2}P^+$, $x_1 + x_2 = 1$, $p_{1,2\perp} = \pm p_\perp$, $q = (0, q^-, q_\perp) =$

$(0, q^-, 0_\perp)$. Then we obtain the form-factors or coupling constants at zero momentum transition[2],

$$
\begin{aligned}
f_{VS\gamma} \;=\;& \frac{e_Q e N_c}{8\pi^3} \int dx_2 d^2 p_\perp \frac{h_V(x_2, p_\perp) h_S(x_2, p_\perp)}{x_1^2 x_2 (M_i^2 - M_0^2)(M_f^2 - M_0^2)} \\
& \left\{ m_Q (M_i^2 - M_f^2) - 2 m_Q x_1 (M_i^2 - M_0^2) - \left[4 m_Q + \frac{x_1 (M_i^2 + M_f^2) + 2 p_\perp^2 - 8 x_1 m_Q^2}{x_1 w_V} \right] p_\perp^2 \right\} \\
& + \frac{e_Q e N_c}{8\pi^3} \int dx_2 d^2 p_\perp \frac{h_V(x_2, p_\perp) h_S(x_2, p_\perp)}{x_1 x_2^2 (M_i^2 - M_0^2)(M_f^2 - M_0^2)} \\
& \left\{ m_Q (M_i^2 - M_f^2) - 2 m_Q x_2 (M_i^2 - M_0^2) - \left[4 m_Q + \frac{x_2 (M_i^2 + M_f^2) + 2 p_\perp^2 - 8 x_2 m_Q^2}{x_2 w_V} \right] p_\perp^2 \right\} ,
\end{aligned} \tag{10}
$$

$$
\begin{aligned}
g_{SV\gamma} \;=\;& \frac{e_Q e N_c}{8\pi^3} \int dx_2 d^2 p_\perp \frac{h_V(x_2, p_\perp) h_S(x_2, p_\perp)}{x_1^2 x_2 (M_i^2 - M_0^2)(M_f^2 - M_0^2)} \\
& \left\{ m_Q (M_f^2 - M_i^2) - 2 m_Q x_1 (M_f^2 - M_0^2) - \left[4 m_Q + \frac{x_1 (M_i^2 + M_f^2) + 2 p_\perp^2 - 8 x_1 m_Q^2}{x_1 w_V} \right] p_\perp^2 \right\} \\
& + \frac{e_Q e N_c}{8\pi^3} \int dx_2 d^2 p_\perp \frac{h_V(x_2, p_\perp) h_S(x_2, p_\perp)}{x_1 x_2^2 (M_i^2 - M_0^2)(M_f^2 - M_0^2)} \\
& \left\{ m_Q (M_f^2 - M_i^2) - 2 m_Q x_2 (M_f^2 - M_0^2) - \left[4 m_Q + \frac{x_2 (M_i^2 + M_f^2) + 2 p_\perp^2 - 8 x_2 m_Q^2}{x_2 w_V} \right] p_\perp^2 \right\} ,
\end{aligned} \tag{11}
$$

where the light-front wave-functions are defined as [12],

$$
\begin{aligned}
h_V(x_2, p_\perp) &= (M_{i/f}^2 - M_0^2) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{\sqrt{2} M_0} \phi_S(x_2, p_\perp) , \\
h_S(x_2, p_\perp) &= (M_{i/f}^2 - M_0^2) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{2\sqrt{6}} \phi_P(x_2, p_\perp) , \\
\phi_P(x_2, p_\perp) &= \sqrt{\frac{2}{\beta^2}} \phi_S(x_2, p_\perp) , \\
\phi_{1S}(x_2, p_\perp) &= 4 \left(\frac{\pi}{\beta^2} \right)^{3/4} \sqrt{\frac{dp_z}{dx_2}} \exp\left(-\frac{p_\perp^2 + p_z^2}{2\beta^2} \right) , \\
\phi_{2S}(x_2, p_\perp) &= 4 \sqrt{\frac{2}{3}} \left(\frac{\pi}{\beta^2} \right)^{3/4} \sqrt{\frac{dp_z}{dx_2}} \exp\left(-\frac{p_\perp^2 + p_z^2}{2\beta^2} \right) \left[\frac{p_\perp^2 + p_z^2}{\beta^2} - \frac{3}{2} \right] , \\
\phi_{3S}(x_2, p_\perp) &= 4 \sqrt{\frac{2}{15}} \left(\frac{\pi}{\beta^2} \right)^{3/4} \sqrt{\frac{dp_z}{dx_2}} \exp\left(-\frac{p_\perp^2 + p_z^2}{2\beta^2} \right) \left[\frac{(p_\perp^2 + p_z^2)^2}{\beta^4} - \frac{5(p_\perp^2 + p_z^2)}{\beta^2} + \frac{15}{4} \right] , \\
\frac{dp_z}{dx_2} &= \frac{e_1 e_2}{x_1 x_2 M_0} , \; w_V = M_0 + 2 m_Q , \\
M_0^2 &= (e_1 + e_2)^2 = \frac{p_\perp^2 + m_Q^2}{x_1} + \frac{p_\perp^2 + m_Q^2}{x_2} , \\
p_z &= \frac{x_2 M_0}{2} - \frac{m_Q^2 + p_\perp^2}{2 x_2 M_0} , \; e_i = \sqrt{m_Q^2 + p_\perp^2 + p_z^2} ,
\end{aligned} \tag{12}
$$

---

[2] For technical details, one can consult Refs.[11, 12].

the M_i and M_f are the masses of the initial and final heavy quarkonium states, the e_i can be viewed as the energy of the quark (antiquark), and the M_0 can be viewed as the kinetic invariant mass of the quark-antiquark system.

In calculations, we have used the following rules [11, 12],

$$\begin{aligned}
\hat{p}_1^\mu &\doteq P^\mu A_1^{(1)} + q^\mu A_2^{(1)} \, , \\
\hat{p}_1^\mu \hat{p}_1^\nu &\doteq g^{\mu\nu} A_1^{(2)} + P^\mu P^\nu A_2^{(2)} + (P^\mu q^\nu + q^\mu P^\nu) A_3^{(2)} + q^\mu q^\nu A_4^{(2)} \, , \\
\hat{N}_2 &\to Z_2 \, , \\
\int d^2 p_\perp \frac{(p_\perp \cdot q_\perp)^2}{q^2} &= -\frac{1}{2} \int d^2 p_\perp \, p_\perp^2 \, , \, \int dx_2 d^2 p_\perp \frac{h_V h_S}{x_2 \hat{N}_1 \hat{N}_1^\prime} \left(x_1 Z_2 - 2 A_1^{(2)} \right) = 0 \, , \\
A_1^{(1)} &= \frac{x_1}{2} \, , \, A_2^{(1)} = \frac{x_1}{2} - \frac{p_\perp \cdot q_\perp}{q^2} \, , \, A_1^{(2)} = -p_\perp^2 - \frac{(p_\perp \cdot q_\perp)^2}{q^2} \, , \\
A_2^{(2)} &= A_1^{(1)} A_1^{(1)} \, , \, A_3^{(2)} = A_1^{(1)} A_2^{(1)} \, , \, A_4^{(2)} = A_2^{(1)} A_2^{(1)} - \frac{A_1^{(2)}}{q^2} \, .
\end{aligned} \tag{13}$$

In the covariant light-front quark models, we usually take the three-dimensional harmonic-oscillator wave-functions in momentum space to approximate the radial wave-functions [12, 15, 18], it is a phenomenological approach. The heavy quarkonium spectra are usually obtained by numerically solving the Schrodinger equation with the Cornell potential or other phenomenological potentials, such as the logarithmic potential, the power-law potential, the QCD-motivated potential [19]. It is difficult to translate the numerical wave-functions into the momentum space and reformulate them with the light-front variables. While the analytical three-dimensional harmonic-oscillator wave-functions are simple, and it is easy to translate them into the momentum space. Furthermore, we cannot solve the light-front hamiltonian as an eigenvalue problem analytically to obtain the light-front wave-functions.

3 Numerical results and discussions

We take the masses of the heavy quarkonium states from the Particle Data Group, $M_{\chi_{c0}} = 3414.75\,\mathrm{MeV}$, $M_{J/\psi} = 3096.916\,\mathrm{MeV}$, $M_{\psi'} = 3686.09\,\mathrm{MeV}$, $M_{\Upsilon} = 9.46030\,\mathrm{GeV}$, $M_{\Upsilon'} = 10.02326\,\mathrm{GeV}$, $M_{\Upsilon''} = 10.3552\,\mathrm{GeV}$, $M_{\chi_{b0}} = 9.85944\,\mathrm{GeV}$, $M_{\chi'_{b0}} = 10.2325\,\mathrm{GeV}$ [6], and obtain the radiative decay widths

$$\begin{aligned}
\Gamma_{\chi_{c0} \to J/\psi \gamma} &= 2214.63 \, e_c^2 e^2 \, g^2_{\chi_{c0} J/\psi \gamma} \, \mathrm{KeV} \, , \\
\Gamma_{\psi' \to \chi_{c0} \gamma} &= 473.535 \, e_c^2 e^2 \, f^2_{\psi' \chi_{c0} \gamma} \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon' \to \chi_{b0} \gamma} &= 113.783 \, e_b^2 e^2 \, f^2_{\Upsilon' \chi_{b0} \gamma} \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon'' \to \chi_{b0} \gamma} &= 3005.49 \, e_b^2 e^2 \, f^2_{\Upsilon'' \chi_{b0} \gamma} \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon'' \to \chi'_{b0} \gamma} &= 48.135 \, e_b^2 e^2 \, f^2_{\Upsilon'' \chi'_{b0} \gamma} \, \mathrm{KeV} \, ,
\end{aligned} \tag{14}$$

Compared to the experimental data from the Particle Data Group [6],

$$\begin{aligned}
\Gamma_{\chi_{c0} \to J/\psi \gamma} &= 122.85 \pm 9.36 \pm 8.4 \; (121.68 \pm 7.02 \pm 8.32) \, \mathrm{KeV} \, , \\
\Gamma_{\psi' \to \chi_{c0} \gamma} &= 27.6848 \pm 1.5488 \pm 0.8866 \; (29.4272 \pm 0.8712 \pm 0.9424) \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon' \to \chi_{b0} \gamma} &= 1.21524 \pm 0.09994 \pm 0.12792 \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon'' \to \chi_{b0} \gamma} &= 0.06096 \pm 0.004995 \pm 0.008128 \, \mathrm{KeV} \, , \\
\Gamma_{\Upsilon'' \to \chi'_{b0} \gamma} &= 1.19888 \pm 0.10915 \pm 0.12192 \, \mathrm{KeV} \, ,
\end{aligned} \tag{15}$$

where the first uncertainties come from the total widths and the second uncertainties come from the branching ratios, we can draw the conclusion tentatively that the coupling constants $f^2_{VS\gamma}$ differ greatly from each other for the bottomonium states, while $g^2_{\chi_{c0} J/\psi \gamma} \approx f^2_{\psi' \chi_{c0} \gamma}$.

From Eqs.(10-12), we can see that the form-factors or coupling constants depend on two kinds of inputs parameters, the constituent quark masses m_Q and the shape parameters β of the wave-functions. The spin averaged ground state masses are $\frac{M_{\eta_c}+3M_{J/\psi}}{4}=3.068\,\mathrm{GeV}$, $\frac{M_{\eta_b}+3M_{\Upsilon}}{4}=9.443\,\mathrm{GeV}$ from the Particle Data Group [6], we estimate the constituent quark masses $m_c\approx 1.5\,\mathrm{GeV}$ and $m_b\approx 4.7\,\mathrm{GeV}$. In this article, we take the constituent quark masses m_Q and shape parameters β of the wave-functions as free parameters and search for the optimal values at the ranges $m_c=(1.3-1.7)\,\mathrm{GeV}$ and $m_b=(4.5-5.0)\,\mathrm{GeV}$.

In numerical calculations, we observe that the form-factors (or coupling constants) are sensitive to the constituent quark masses m_Q and the shape parameters β, small variations of those input parameters can lead to large changes of the form-factors (thereafter the decay widths); the optimal values are $m_c=1.5\,\mathrm{GeV}$ and $m_b=4.8\,\mathrm{GeV}$, $\beta_{\chi_{c0}}=1.30\,\mathrm{GeV}$, $\beta_{J/\psi}=0.76\,\mathrm{GeV}$, $\beta_{\psi'}=0.83\,\mathrm{GeV}$, $\beta_{\Upsilon'}=1.30\,\mathrm{GeV}$, $\beta_{\Upsilon''}=0.976\,\mathrm{GeV}$, $\beta_{\chi_{b0}}=0.94\,\mathrm{GeV}$, $\beta_{\chi'_{b0}}=0.945\,\mathrm{GeV}$, which can reproduce the decay widths of the five observed processes $\chi_{c0}\to J/\psi\gamma$, $\psi'\to\chi_{c0}\gamma$, $\Upsilon'\to\chi_{b0}\gamma$, $\Upsilon''\to\chi_{b0}\gamma$ and $\Upsilon''\to\chi'_{b0}\gamma$ [6]. If the form-factors (or coupling constants) are not sensitive to the shape parameters β of the wave-functions, we can introduce two parameters to characterize the charmonium and bottomonium states respectively, and search for the optimal values to reproduce the decay widths of the five observed radiative transitions, however, two parameters cannot lead to satisfactory results.

The optimal values $m_c=1.5\,\mathrm{GeV}$ and $m_b=4.8\,\mathrm{GeV}$ approximate or equal to the estimated values $m_c\approx 1.5\,\mathrm{GeV}$ and $m_b\approx 4.7\,\mathrm{GeV}$. In the constituent quark model, the usually used constituent quark masses are $m_c=(1.4-1.6)\,\mathrm{GeV}$ and $m_b=(4.7-4.9)\,\mathrm{GeV}$, it is natural to take the values $m_c=1.5\pm 0.1\,\mathrm{GeV}$ and $m_b=4.8\pm 0.1\,\mathrm{GeV}$.

We choose (or suppose) the shape parameters have the same uncertainties as the constituent quark masses tentatively, i.e. $\beta_{\chi_{c0}}=1.30\pm 0.1\,\mathrm{GeV}$, $\beta_{J/\psi}=0.76\pm 0.1\,\mathrm{GeV}$, $\beta_{\psi'}=0.83\pm 0.1\,\mathrm{GeV}$, $\beta_{\Upsilon'}=1.30\pm 0.1\,\mathrm{GeV}$, $\beta_{\Upsilon''}=0.976\pm 0.1\,\mathrm{GeV}$, $\beta_{\chi_{b0}}=0.94\pm 0.1\,\mathrm{GeV}$, $\beta_{\chi'_{b0}}=0.945\pm 0.1\,\mathrm{GeV}$, and study the resulting uncertainties,

$$\begin{aligned}
\Gamma_{\chi_{c0}\to J/\psi\gamma} &= 121.54^{+160.22\,+34.32\,+39.89}_{-97.18\,-26.26\,-32.67}\,\mathrm{KeV}\,,\\
\Gamma_{\psi'\to\chi_{c0}\gamma} &= 27.90^{+26.67\,+9.83\,+1.56}_{-17.97\,-9.10\,-3.55}\,\mathrm{KeV}\,,\\
\Gamma_{\Upsilon'\to\chi_{b0}\gamma} &= 1.23^{+2.31\,+0.07\,+0.52}_{-1.13\,-0.13\,-0.34}\,\mathrm{KeV}\,,\\
\Gamma_{\Upsilon''\to\chi_{b0}\gamma} &= 0.0606^{+0.0078\,+0.2160\,+0.2343}_{-0.0072\,-0.0605\,-0.0589}\,\mathrm{KeV}\,,\\
\Gamma_{\Upsilon''\to\chi'_{b0}\gamma} &= 1.19^{+0.31\,+2.30\,+3.75}_{-0.27\,-1.18\,-1.14}\,\mathrm{KeV}\,,
\end{aligned}\tag{16}$$

where the uncertainties come from the heavy quark masses and shape parameters of the initial and final quarkonium states, sequentially.

From Eqs.(15-16), we can see that the central values of the present predictions are consistent with the experimental data from the Particle Data Group [6], however, the uncertainties come from the heavy quark masses exceed 100% for the transitions $\chi_{c0}\to J/\psi\gamma$, $\psi'\to\chi_{c0}\gamma$, $\Upsilon'\to\chi_{b0}\gamma$, and the uncertainties come from the shape parameters also exceed 100% for the transitions $\Upsilon''\to\chi_{b0}\gamma$, $\Upsilon''\to\chi'_{b0}\gamma$. We have little room for varying those parameters to reproduce the experimental data, which weakens the predictive ability of the covariant light-front quark model remarkably. On the other hand, if we take the uncertainties of the experimental data on the radiative transitions (see Eq.(15)) to constrain the uncertainties of the constituent quark masses and shape parameters, the allowed uncertainties are $|\delta m_c|<10\,\mathrm{MeV}$, $|\delta m_b|<10\,\mathrm{MeV}$, $|\delta\beta_{\chi_{c0}}|<35\,\mathrm{MeV}$, $|\delta\beta_{J/\psi}|<30\,\mathrm{MeV}$, $|\delta\beta_{\psi'}|<17\,\mathrm{MeV}$, $|\delta\beta_{\Upsilon'}|<100\,\mathrm{MeV}$, $|\delta\beta_{\Upsilon''}|<8\,\mathrm{MeV}$, $|\delta\beta_{\chi_{b0}}|<10\,\mathrm{MeV}$, $|\delta\beta_{\chi'_{b0}}|<6\,\mathrm{MeV}$.

The heavy quarkonium states have equal constituent quark masses, $m_Q=m_{\overline{Q}}$, the heavy quark masses are taken as one parameter m_Q rather than two parameters m_Q and $m_{\overline{Q}}$, see Eqs.(10-12), the total uncertainties come from the heavy quark masses are larger than that come from the heavy quark masses m_Q and $m_{\overline{Q}}$, respectively.

In Figs.2-3, we plot the radiative decay widths $\Gamma_{\chi_{c0}\to J/\psi\gamma}$, $\Gamma_{\psi'\to\chi_{c0}\gamma}$, $\Gamma_{\Upsilon'\to\chi_{b0}\gamma}$, $\Gamma_{\Upsilon''\to\chi_{b0}\gamma}$ and $\Gamma_{\Upsilon''\to\chi'_{b0}\gamma}$ with variations of the shape parameters β. From the figures, we can see that the widths are sensitive to the shape parameters β indeed, small variations of the shape parameters β can lead to rather large changes of the decay widths, especially when there are nodi in the radial wave-functions, it is very difficult (or impossible) to obtain two universal parameters $\beta_{\bar{c}c}$ and $\beta_{\bar{b}b}$ to characterize the charmonium and bottomonium states respectively.

We can take the optimal values $m_c = 1.5\,\text{GeV}$ and $m_b = 4.8\,\text{GeV}$, $\beta_{\chi_{c0}} = 1.30\,\text{GeV}$, $\beta_{J/\psi} = 0.76\,\text{GeV}$, $\beta_{\psi'} = 0.83\,\text{GeV}$, $\beta_{\Upsilon'} = 1.30\,\text{GeV}$, $\beta_{\Upsilon''} = 0.976\,\text{GeV}$, $\beta_{\chi_{b0}} = 0.94\,\text{GeV}$, $\beta_{\chi'_{b0}} = 0.945\,\text{GeV}$ as the basic input parameters and study other radiative transitions among the heavy quarkonium states. For example, we can take the parameter $\beta_{\chi_{b0}} = 0.94\,\text{GeV}$ to calculate the decay width $\Gamma_{\chi_{b0}\to\Upsilon\gamma}$. From Fig.3, we can see that the prediction $\Gamma_{\chi_{b0}\to\Upsilon\gamma} = (18-34)\,\text{KeV}$ for $\beta_{\Upsilon} = (0.4-1.4)\,\text{GeV}$ is consistent with the value $\Gamma_{\chi_{b0}\to\Upsilon\gamma} = (24-26)\,\text{KeV}$ from the potential quark models [20]. If we take the parameter $\beta_{\Upsilon} = 1.16\,\text{GeV}$, the predictions $\Gamma_{\chi_{b0}\to\Upsilon\gamma} = 24.75\,\text{KeV}$ and $\Gamma_{\chi'_{b0}\to\Upsilon\gamma} = 5.97\,\text{KeV}$ are consistent with the values from the potential quark models $\Gamma_{\chi_{b0}\to\Upsilon\gamma} = (24-26)\,\text{KeV}$ and $\Gamma_{\chi'_{b0}\to\Upsilon\gamma} = (4.5-8.5)\,\text{KeV}$ [20], while the prediction $\Gamma_{\chi'_{b0}\to\Upsilon'\gamma} = 46.74\,\text{KeV}$ is much larger than the value $\Gamma_{\chi'_{b0}\to\Upsilon'\gamma} = (11-12)\,\text{KeV}$ from the potential quark model [20].

Without precise experimental data, we cannot determine the value of the shape parameter β_{Υ}. On the other hand, there are controversies for the spectroscopy of the charmonium states, and lack experimental data on other radiative transitions among the vector and scalar charmonium states. We prefer study those processes in the future.

In Ref.[15], W. Wang calculates the S-wave to S-wave radiative transitions $\Upsilon \to \eta_b\gamma$, $\Upsilon' \to \eta_b\gamma$, $\Upsilon' \to \eta'_b\gamma$, $\Upsilon'' \to \eta_b\gamma$, $\Upsilon'' \to \eta'_b\gamma$, $J/\psi \to \eta_c\gamma$, $\psi' \to \eta_c\gamma$, $\psi' \to \eta'_c\gamma$ with the covariant light-front quark model, and observes that those M_1 transitions are sensitive to the heavy quark masses and shape parameters of the wave-functions, the existing experimental data cannot be reproduced consistently with suitable parameters. The input parameters used in studying the observed M_1 transitions again fail to reproduce the experimental data on the E_1 transitions of the heavy quarkonium states. In Ref.[21], H. W. Ke et al observe that the three-dimensional harmonic-oscillator wave-functions lead to obvious contradictions between the experimental and theoretical values of the $\Upsilon(nS)$ decay constants. They modify the wave-functions by introducing several additional parameters to describe the radial quantum number n dependence and keep the orthogonality among the nS states, then fit the decay constants to determine the $\Upsilon(nS)$ wave-functions and study the radiative decays $\Upsilon(nS) \to \eta_b + \gamma$, the prediction of the $\Upsilon(3S) \to \eta_b + \gamma$ is in accordance with the experimental data by the order of magnitude, the discrepancy between the experimental and theoretical values is not well smeared. In the present case, the different values of the shape parameters $\beta_{\chi_{c0}}$, $\beta_{J/\psi}$, $\beta_{\psi'}$, $\beta_{\Upsilon'}$, $\beta_{\Upsilon''}$, $\beta_{\chi_{b0}}$, $\beta_{\chi'_{b0}}$ violate the orthogonality of the wave-functions, it is the shortcoming of the present work. We can also draw the conclusion tentatively that we cannot get rid of the sensitivity to the shape parameters β without introducing additional parameters in the present case, however, the experimental data are far from enough to fit the additional parameters.

4 Conclusions

In this article, we study the radiative transitions among the vector and scalar heavy quarkonium states in the framework of the covariant light-front quark model. In calculations, we observe that the radiative decay widths are sensitive to the constituent quark masses and the shape parameters of the light-front wave-functions. We reproduce the experimental data for the observed processes with suitable parameters, while the predictions for the un-observed processes are consistent or inconsistent with other theoretical calculations. The parameters can be fitted to the precise experimental data in the future, although it is a difficult work.



Figure 2: The radiative decay widths with variations of the shape parameter $\beta_{\chi_{c0}}$. In (I) $\chi_{c0} \to J/\psi\gamma$, A, B, C, D correspond to $\beta_{J/\psi}/\beta_{\chi_{c0}} = 0.56/1.3$, $0.66/1.3$, $0.76/1.3$, $0.86/1.3$, respectively; in (II) $\psi' \to \chi_{c0}\gamma$, A, B, C, D correspond to $\beta_{\psi'}/\beta_{\chi_{c0}} = 0.63/1.3$, $0.73/1.3$, $0.83/1.3$, $0.93/1.3$, respectively.



Figure 3: The radiative decay widths with variations of the shape parameters $\beta_{\chi_{b0}}$ or $\beta_{\chi'_{b0}}$. In (I) $\Upsilon' \to \chi_{b0}\gamma$, A, B, C, D correspond to $\beta_{\Upsilon'}/\beta_{\chi_{b0}} = 1.1/0.94$, $1.2/0.94$, $1.3/0.94$, $1.4/0.94$, respectively; in (II) $\Upsilon'' \to \chi_{b0}\gamma$, A, B, C, D correspond to $\beta_{\Upsilon''}/\beta_{\chi_{b0}} = 0.776/0.94$, $0.876/0.94$, $0.976/0.94$, $1.076/0.94$, respectively; in (III) $\Upsilon'' \to \chi'_{b0}\gamma$, A, B, C, D correspond to $\beta_{\Upsilon''}/\beta_{\chi'_{b0}} = 0.776/0.945$, $0.876/0.945$, $0.976/0.945$, $1.076/0.945$, respectively; in (IV) $\chi_{b0} \to \Upsilon\gamma$, $\beta_{\Upsilon} = 0.94\,$GeV.

Acknowledgment

This work is supported by National Natural Science Foundation of China, Grant Number 11075053, and the Fundamental Research Funds for the Central Universities.

References

[1] M. Ablikim et al, Phys. Rev. Lett. **109** (2012) 172002.

[2] N. E. Adam et al, Phys. Rev. Lett. **94** (2005) 232002.

[3] H. Mendez et al, Phys. Rev. **D78** (2008) 011102.

[4] Z. G. He, X. R. Lu, J. Soto and Y. Zheng, Phys. Rev. **D83** (2011) 054028.

[5] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, F. Feruglio, R. Gatto and G. Nardulli, Phys. Rept. **281** (1997) 145.

[6] J. Beringer et al, Phys. Rev. **D86** (2012) 010001.

[7] H. Grotch, D. A. Owen and K. J. Sebastian, Phys. Rev. **D30** (1984) 1924; W. Kwong and J. L. Rosner, Phys. Rev. **D38** (1988) 279; M. B. Voloshin, Prog. Part. Nucl. Phys. **61** (2008) 455; E. Eichten, S. Godfrey, H. Mahlke and J. L. Rosner, Rev. Mod. Phys. **80** (2008) 1161.

[8] J. J. Dudek, R. G. Edwards and D. G. Richards, Phys. Rev. **D73** (2006) 074507; J. J. Dudek, R. Edwards and C. E. Thomas, Phys. Rev. **D79** (2009) 094504; Y. Chen et al, Phys. Rev. **D84** (2011) 034503;

[9] N. Brambilla, P. Pietrulewicz and A. Vairo, Phys. Rev. **D85** (2012) 094005.

[10] N. Brambilla et al, Eur. Phys. J. **C71** (2011) 1534.

[11] W. Jaus, Phys. Rev. **D60** (1999) 054026.

[12] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. **D69** (2004) 074025.

[13] H. Y. Cheng and C. K. Chua, Phys. Rev. **D69** (2004) 094007.

[14] H. M. Choi, C. R. Ji and L. S. Kisslinger, Phys. Rev. **D65** (2002) 074032; C. W. Hwang and Z. T. Wei, J. Phys. **G34** (2007) 687; H. M. Choi, Phys. Rev. **D75** (2007) 073016; W. Wang, Y. L. Shen and C. D. Lu, Eur. Phys. J. **C51** (2007) 841; W. Wang, Y. L. Shen and C. D. Lu, Phys. Rev. **D79** (2009) 054012; H. Y. Cheng and C. K. Chua, Phys. Rev. **D81** (2010) 114006; T. Peng and B. Q. Ma, Eur. Phys. J. **A48** (2012) 66.

[15] W. Wang, arXiv:1002.3579.

[16] H. W. Ke, X. Q. Li and X. Liu, arXiv:1002.1187.

[17] G. Bonvicini et al, Phys. Rev. **D70** (2004) 032001.

[18] C. W. Hwang, Eur. Phys. J. **C62** (2009) 499.

[19] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane and T. M. Yan, Phys. Rev. **D17** (1978) 3090; E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane and T. M. Yan, Phys. Rev. **D21** (1980) 203; E. J. Eichten and C. Quigg, Phys. Rev. **D49** (1994) 5845.

[20] B. Q. Li and K. T. Chao, Commun. Theor. Phys. **52** (2009) 653; and references therein.

[21] H. W. Ke, X. Q. Li, Z. T. Wei and X. Liu, Phys. Rev. **D82** (2010) 034023.